



For Healthy Liver

# BIKARIN

Cap./Susp.

BKR

Each capsule contains **Silymarin 70mg** with **Lecithin 275 mg** + B-Complex  
Each 5ml contains **Silymarin 35mg** with **Lecithin 275 mg**

## LIVER PROTECTOR

- ◆ Prevents entry of toxins in liver cells
- ◆ Treatment of liver cirrhosis by Vit.B
- ◆ Regulates fat metabolism
- ◆ Stabilizes cell membrane
- ◆ Non Alcoholic Fatty Liver Disease
- ◆ Drug Induced Liver Damage
- ◆ Non Alcoholic Liver Cirrhosis

### DOSE

Children : 2.5 ml BID  
Adults : 5-10 ml BID  
1-2 Cap. BID



# BIKARIN-L

Cap./Susp.

Each capsule contains **Silymarin 140mg** with Lecithin 525mg  
Each 5ml contains **Silymarin 70mg** with Lecithin 275mg

BKR-L

## For Treatment of Liver Diseases

### Silymarin

- ◆ Proven antioxidant
- ◆ Stabilizes Hypatocyte Membrane, regulates permeability.
- ◆ Stimulates ribosomal RNA Synthesis promoting Liver cell regeneration.
- ◆ Prevents transformation of stellate hepatocytes into myofibroblast which is responsible for deposition of Collagen fibre resulting cirrhosis.  
– Pharmacology of silymarin, Clinical Drug Investigation 22(1):51-65 2002
- ◆ Silymarin has hepatoprotectant properties, it has been used in Various Liver disorders as well as to prevent hepatotoxicity.  
– Martindale, 36th Ed. P:1452

- ◆ Alcoholic Hepatitis
- ◆ Alcoholic Fatty Liver
- ◆ Alcoholic Liver Cirrhosis



### DOSE

Children : 5 ml BID  
Adults : 10 ml BID  
1 Cap. BID

### **Rx** Bikarin-L Suspension

- Jaundice
- Alcoholic Liver Diseases
- Viral Hepatitis
- Liver cirrhosis
- NAFLD
- TOXIC or Drug induced Liver disease

**Healthy & active life is the secret of happy life**  
**ENSURE HEALTHY & ACTIVE LIFE TO ALL**



**Rx Bikaton-L syrup**

- « Lack of appetite
- « Convalescence
- « General debility
- « Growing children
- « Underweight infants and children

*with*  
**Bikaton-L**  
**Syrup**

Each 5ml contains L-Lysine fortified 375mg with 9mg Betacarotene Zinc, B-Complex and Minerals.

BKT


Each 5ml offers	
L-Lysine Monohydrate	375 mg.
Beta-Carotene (2.5% dispersion)	9 mg.
Eqv. to Vit. A	2500 I.U.
Vit. D3	200 I.U.
Benfotiamine (Vit B <sub>1</sub> )	2.5 mg.
Pyridoxin Hcl (Vit B <sub>6</sub> )	0.75 mg.
Methylcobalamin	50 mcg.
Vit. B <sub>2</sub>	1 mg.
Niacinamide	5 mg.
Biotin	50 mcg.
Zinc Sulphate Monohydrate	
Eqv. to elemental Zinc	8.8 mg.
Chromium Picolinate	
Eqv. to elemental Chromium	30 mcg.
Sodium Selenate	
Eqv. to elemental Selenium	10 mcg.
Inositol	10 mg.

- « Regulates metabolic function
- « Improves appetite
- « Restores healthy & active life to infants, children and adults

L-Lysine helps to increase appetite by detoxifying liver & by helping to produce some enzymes that help digestion :

– Clinical Nutrition 1992

**DOSE**  
 Children : 5 ml BID  
 Adults : 10 ml BID



# Bikaton-L

Softgel Capsule

BKT  
GEL

**Omega 3 Fatty Acids**  
(EPA AND DHA)

+

**6 Natural Extracts**  
(Ginseng / Garlic / Green Tea / Gingko Biloba /  
Grape Seed / Wheat Germ oil)

+

**11 Minerals**  
(Ca + Cu + Cr + Bo + I + Fe + Mg + Mn + K + Zn +Se)

+

**10 Vitamins**  
(A + B + B<sub>3</sub> + B<sub>6</sub> + B<sub>7</sub> + B<sub>9</sub> + B<sub>12</sub> + C + D<sub>3</sub> + K)

**Bring Back The "Golden Years" of Youth**





# NODOREX

Exp.

Each 5ml contains Levocetirizine 2.5mg + Ambroxol 15mg + Guaiphenesin 50mg + Menthol 5mg

## Levocetirizine:

Reduces cough intensity and frequency<sup>1</sup>

## Ambroxol:

Promotes mucus clearance<sup>2</sup>

## Guaiphenesin:

Decreases sputum thickness in 96% patients<sup>3</sup>

## Menthol:

Offers soothing effect


- » Liquifies Viscid sputum
- » Soothes Irritated Throat
- » Helps expectoration
- » Restores soothing comfort

- DAY TO DAY COUGH
- ALLERGIC COUGH
- SEASONAL COUGH & COLD
- COUGH OF LRTI,URTI

### DOSE

Children : 5 ml BID  
Adults : 10 ml BID

NOD  
EXP





# NODOREX-DX

Syrup

Each 5ml contains Levocetirizine 2.5mg + Dextromethorphan 10mg + Methol 2.5mg

## Levocetirizine:

Reduces cough intensity and frequency<sup>1</sup>

## Dextromethorphan

Lowers cough intensity to a greater degree than codeince<sup>2</sup>

## Guaiphenesin:

Decreases sputum thickness in 96% patients<sup>3</sup>

## Menthol:

Offers soothing effect

- » Calms cough center
- » Relieves dry & irritating Cough
- » Brings back soothing solution.

- HACKING COUGH
- TB COUGH
- SMOKER'S COUGH
- DRY & IRRITATING COUGH


### DOSE

Children : 5 ml BID  
Adults : 10 ml BID

NOD  
DX



When **coughing** bothers daily life, for soothing solution



# NODOREX-LS

Syrup

Each 5ml contains Levosalbutamol 1mg + Ambroxol 30mg + Guaiphenesin 50mg

## Levosalbutamol


- ❖ 4 times more potent than racaemic Salbutamol<sup>2</sup>
- ❖ Offers better safety profile and lesser side-effects than Salbutamol<sup>2</sup>
- ❖ More effective than Salbutamol in terms of Asthma score and PEF<sup>3</sup>

NOD  
LS

**Indicated in** Cough associated with **Asthma, Bronchitis and COPD**

### DOSE

Adults : 5-10 ml BID





# NODOREX-LS

*Junior*

Each 5ml contains Levosalbutamol 0.5mg + Ambroxol 15mg + Guaiphenesin 50mg

## Levosalbutamol

- ✦ Efficacy and Safety in neonates is **endorsed by the Indian Journal of Paediatrics**<sup>4</sup>

## Ambroxol

- ✦ Offers mukokinetic and mucolytic action and found **effective in 92% patients**<sup>5</sup>


## Guaiphenesin

- ✦ **Powerful expectorant**, reduces mucus thickness to facilitate expectoration<sup>6</sup>

**Indicated in** Cough associated with **Asthma, Bronchitis and COPD**

### DOSE

Children :(3-12 yrs)  
5-10 ml BID





**IN ACUTE  
WATERY  
DIARRHOEA**



Diarrhoea cause nearly half million deaths of Indian children under the age of five years. Majority of deaths are due to hypersecretion of water & electrolytes resulting dehydration.

-United State Agency for International Development (USAID). March 2010.

# MOTIRIL<sup>Susp.</sup>

Each 5ml contains Racecadotril 15mg

Racecadotril inhibits endogenous enkephalinase, restores intestinal enkephalins, inhibits hypersecretion of water & electrolytes, **prevent dehydration.**

- « Safe & effective treatment to control hypersecretion in diarrhoea
- « Effectively prevents dehydration
- « Easy to titrate dose

**Dose:**

Infants & Children  
1.5mg/kg. Every 8hours till  
Diarrhoea controlled.

**Age wise dose (approx)**

Up to 1year age :1tsf TID  
Above 1 year up to 2 years: 1-1.5tsf TID  
Above 2 year up to 3 years: 1.5-2tsf TID  
Above 3 year up to 6 years: 2tsf TID

"Enkephalins, the enteric neurotransmitter, inhibit intestinal secretion. Enkephalinase, an endogenous enzyme, inhibits enkephalins, resulting hypersecretion Racecadotril inhibits enkephalinase, restores enkephalins, produce antidiarrhoeal effect"

– Goodman & Gilman's  
11th Ed. P: 997

# ZINCONAK

SYRUP

Each 5ml contains Zinc Acetate eq. to Elemental Zinc 20mg

- ◆ Shortens the average duration of diarrhoea by 16 hours in Children<sup>1</sup>
- ◆ Zinc deficiency may have adverse effects on physical growth and neurodevelopment<sup>2</sup>
- ◆ Reduce the frequency and severity of respiratory infections in children<sup>3</sup>

## INDICATED IN

- ◆ Persistent Diarrhoea
- ◆ Growth & Development
- ◆ Boosting Immunity

## USAGE

Syrup : 5 ml OD



ZIN

*Get More out of your preferred Zinc*

# OILAED *Oil*

Vit. A 20,000IU + Vit. D 4000IU + Vit. E 4.8mg + Olive Oil 10% / per 10ml

## OFFERS 10% MORE OLIVE OIL

- ◆ Olive Oil massage **improves health** and weight gain in Newborns<sup>1</sup>
- ◆ Offers **10% extra virgin olive oil**, best carrier of Vitamins
- ◆ Infants who receive regular oil massage sleep better and cry less than others<sup>2</sup>
- ◆ Vitamin D supplementation in childhood is **endorsed by WHO and UNICEF** for healthy bones and baby.

### USAGE

A gentle massage in newborns and infants before bath



OAD  
OIL

# Safer Sedation For Infants & Children



- Insomnia
  - Short term sleeping problem
  - Restlessness
  - Vaccination
- Infants upto 1 yr. of age: 20-30mg/kg 30min. before bedtime  
Children upto 6 yrs. of age: 30-50mg/kg 30 min. before bedtime

- CT- Scan
  - MRI
  - Dressing/I.V.Lines
  - Other Surgical Procedures
- Children upto 6 yrs. of age: 50-75mg/kg, 30-60 min. before procedure

# KIDPOSED DROP SYRUP

Each 1ml Drop : Triclofos Sodium 100mg  
Each 5ml Syrup: Triclofos Sodium 500mg

## DISTINCTLY SUPERIOR TO CHLORAL HYDRATE

- ▣ Provides early onset of sleep up to median duration of 90 mins<sup>1</sup>
  - ▣ Triclofos is an effective sedative in children<sup>2</sup>
  - ▣ Offers better palatability and less gastric irritation as compared to chloral hydrate<sup>2</sup>
- ♦ Shortens the time of onset of sleep
  - ♦ Ensures 6-8 hours sleep
  - ♦ No hangover
  - ♦ High safety margin

*"Triclofos is stable ester of Trichloroethanol, The derivative of chloral hydrate. It is specially widely employed in children and in elderly people because it is well tolerated and is a safer drug."*

*- Clinical Pharmacology  
D.R.Laurence, 4th Ed. p:10.7*

**DOSE**  
30 to 50 mg/kg B.W. 45-60 minutes before procedure.\*  
(Max. up to 100 mg/kg and Max. of 2 gm)



In Abdominal discomfort | Dyspepsia | Indigestion

# ZYMOAID

Fungal Diastase 50mg + Pepsin 10mg in each 5ml

Syrup



The *tasty* Digestive with *Sorbitol* Advantage




PEPSIN - Digests Protein

FUNGAL DIASTASE - Digests carbohydrate

SORBITOL - Osmotic laxative that relieves constipation

#### DOSE

5 - 10ml  
after each meal



ZYM

TIME TESTED PROBIOTIC THAT PROTECTS  
INTESTINAL FLORA IN INFANTS & CHILDREN



**Rx** **Esbiped Sachet**

- ▶ Antibiotic Associated Diarrhoea
- ▶ Alongwith Oral Antibiotics

**Usual Dose :** 1-2 sachets twice daily

# Esbiped **SACHET**

Each Sachet offers

**Saccharomyces Boulardii 282.5mg**  
(Corresponding to yeast 250mg)

**TO RESTORE INTESTINAL FLORA IN INFANTS & CHILDREN**

- ◆ Combats c.difficile,v.cholerae & E.coli toxins
- ◆ Restores microbalance in colon and small Intestine
- ◆ Superior to lactobacilli, which act only in colon

“SaccharomycesBoulardii has demonstrated activity In colon and small intestine in contrast to Lactobacillus which has effect in colon only;”

-Digestion 1996.20:192-196.

**DOSE**

1-2 Sachet BID



ESB

Bacterial or Mixed Diarrhea  
The dependable treatment for Children & Adults.



For treatment of diarrhea

<sup>Rx</sup> **Harlox-OZ** SUSPENION 1 tsf twice a day for Children  
<sup>Rx</sup> **Harlox-OZ** TABLET 1 tab twice a day for Adults.

# Harlox-OZ

Susp.

Each 5ml contains Ofloxacin 50mg with Ornidazole 125mg

# Harlox-OZ

Tablet

Each tablet contains Ofloxacin 200mg with Ornidazole 500mg

● **Ofloxacin**

Timetested antibacterial for G.I. disorders

● **Ornidazole**

The broadspectrum antiprotozoal



HAR

Damage to myelin sheath is the most obvious lesion in neuropathy. Evidences suggest that impairment of methionine synthetase reaction in deficiency of methylcobalamin is likely to be responsible for this abnormality.

- Goodman & Gilman's, 10th ed. p: 1505




R<sub>x</sub> **NCOB** Capsule

- « Peripheral neuropathy
- « Diabetic neuropathy
- « Alcoholic neuropathy

- « Burning sensation
- « Muscle cramps
- « Neuralgia

AIDONAK Offers

# NCOB Capsule



Each CAPSULE contains Methylcobalamin 1500meg  
Vit.B6 3mg, Folic Acid 1.5mg, Alpha-lipoic Acid 100mg

## Methylcobalamin

- ◆ Essential cofactor for Methionine synthetase reaction.
- ◆ Converts homocysteine to Methionine.
- ◆ Methionine is essential for Synthesis of myelin basic protein.
- ◆ Prevents & repair damage to myelin sheath, rejuvenates neurons.

## Alpha-lipoic Acid

- ◆ Potent antioxidant
- ◆ Regenerates Vitamin E
- ◆ Prevents diabetic polyneuropathy
- ◆ Helps improve nerve function in diabetic Patients

— Diabetic Care '1999.

### DOSE

1 capsule once or twice daily  
for 30-90 days or more



NCOB





When the objective is prevention of iron deficiency in pregnant women, an amount of 15mg to 30mg iron per day are adequate to meet the requirement in 2nd & 3rd trimester"

– Harrison's, 15th Ed. Vol.1.

***Rx* HEMGARD-XT SYRUP**

Usual dose: 1-2 tsf daily

***Rx* HEMGARD-XT TABLET**

Usual dose: 1 Tablet OD

B12, Folic acid, Iron is the need in Therapy of Oral Iron

# HEMGARD-XT

Each Tablet contains Ferrous Ascorbate Eqv. to elemental Iron 100mg + Folic Acid 1.5mg

Each 5ml Syrup contains Ferrous Ascorbate Eqv. to elemental Iron 30mg + Folic Acid 0.55mg + Vitamin B12 7.5mcg

## Ferrous Ascorbate

- ◆ Better iron absorption
- ◆ No constipation / G.I. complications
- ◆ Assured hemoglobin response

"Combination of Ferrous compound with reducing sugars or ascorbate prevents or retards oxidation of Ferrous molecule resulting an increased absorption of iron. Ascorbate prevents oxidation of ferrous iron, and Ferrous ascorbate release ferrous ions directly to gastric mucosa from ascorbate complex."

– Dept. Of Pharmaceutical Analysis & Analytical Chemistry. University of Leiden, Netherlands



HEM



- Dental Pain
- Headache
- Fever

- Fracture & Dislocations
- Dysmenorrhoeal Pain
- Sprain & Strain
- Low back Pain,
- Arthritic Pain



# AIDOFEN-P

Tablet

Each Tablet contains Aceclofenac 100mg + Paracetamol 325mg

# AIDOFEN-SP

Tablet

Each Tablet contains Aceclofenac 100mg + Serratiopeptidase 15mg + Paracetamol 325mg

Effective treatment for day to day patient care for Pain, Inflammation & Fever.

- Site Specific Action
- Prompt Pain Relief

### DOSE

1 Tablet Twice a day



# STAYRAB-DSR

Capsule

Rabeprazole 20mg. enteric coated granules  
Domperidone 30mg. sustained release granules

## **RABEPRAZOLE**

- ◆ Proven proton pump inhibitor, ensures round the Clock acid inhibition.

## **DOMPERIDONE**

- ◆ Time tested prokinetic agent.
- ◆ Restores normal motility & tone of upper GIT.
- ◆ Facilitates gastric emptying.

## **<sup>Rx</sup> STAYRAB-DSR Capsule**

- ◆ GERD
- ◆ Epigastric sense of fullness
- ◆ Heartburn
- ◆ Nausia & Vomiting
- ◆ With NSAIDS
- ◆ Diabetic gastroparesis.

### **DOSE**

1 Tablet a day



The ideal tieup between Acid inhibitory and Prokinetic agent

DEPENDABLE DERMACARE IN

# BIKADERM

## Cream

Clobetasol Propionate 0.05% (w/w) + Neomycin Sulphate 0.05% (w/w) + Miconazole Nitrate 2% (w/w)

### Clobetasol:

- ◆ Potent topical anti-inflammatory

### Neomycin:

- ◆ Potent topical anti-fungal

### Miconazole:

- ◆ Potent topical anti-bacterial

*Easy washable  
Nonstaining to clothes*

### INDICATED IN

- ◆ Contact Dermatitis
- ◆ Candida and Tinea Infections
- ◆ Eczema

In Mixed Skin Infections  
Choose the "All in One" formula

Usually two applications a day

