

**IN ACUTE
WATERY
DIARRHOEA**



Diarrhoea cause nearly half million deaths of Indian children under the age of five years. Majority of deaths are due to hypersecretion of water & electrolytes resulting dehydration.

-United State Agency for International Development (USAID). March 2010.

MOTIRIL^{Susp.}

Each 5ml contains Racecadotril 15mg

Racecadotril inhibits endogenous enkephalinase, restores intestinal enkephalins, inhibits hypersecretion of water & electrolytes, **prevent dehydration.**

- « Safe & effective treatment to control hypersecretion in diarrhoea
- « Effectively prevents dehydration
- « Easy to titrate dose

Dose:

Infants & Children
1.5mg/kg. Every 8hours till
Diarrhoea controlled.

Age wise dose (approx)

Up to 1year age :1tsf TID
Above 1 year up to 2 years: 1-1.5tsf TID
Above 2 year up to 3 years: 1.5-2tsf TID
Above 3 year up to 6 years: 2tsf TID

"Enkephalins, the enteric neurotransmitter, inhibit intestinal secretion. Enkephalinase, an endogenous enzyme, inhibits enkephalins, resulting hypersecretion Racecadotril inhibits enkephalinase, restores enkephalins, produce antidiarrhoeal effect"

– Goodman & Gilman's
11th Ed. P: 997